

ASYMMETRIC TRUNCATED TOEPLITZ OPERATORS ON FINITE-DIMENSIONAL SPACES II

BARTOSZ ŁANUCHA

ABSTRACT. In this paper we present some consequences of the description of matrix representations of asymmetric truncated Toeplitz operators acting between finite-dimensional model spaces. In particular, we prove that these operators can be characterized using modified compressed shifts or the notion of shift invariance. We also describe rank-one asymmetric truncated Toeplitz operators acting between finite-dimensional model spaces. This paper is a sequel to the paper of J. Jurasik and B. Łanucha *Asymmetric truncated Toeplitz operators on finite-dimensional spaces*.

1. INTRODUCTION

Let H^2 be the classical Hardy space of the unit disk $\mathbb{D} = \{z: |z| < 1\}$ and let P denote the Szegő projection, that is, the orthogonal projection from $L^2(\partial\mathbb{D})$ onto H^2 .

The unilateral shift $S: H^2 \rightarrow H^2$ is defined on H^2 by

$$Sf(z) = zf(z).$$

It is known that the S -invariant subspaces of H^2 can be characterized in term of inner functions. An inner function α is a function from H^∞ , the algebra of bounded analytic functions on $\mathbb{D}$, such that $|\alpha| = 1$ a.e. on $\partial\mathbb{D}$. The theorem of A. Beurling states that every closed nontrivial S -invariant subspace of H^2 is of the form αH^2 for some inner function α . Consequently, each of the closed nontrivial subspaces of H^2 invariant under the backward shift S^* is given by

$$K_\alpha = H^2 \ominus \alpha H^2$$

with α inner. The space K_α is called the model space corresponding to the inner function α .

The model space K_α has a reproducing kernel property and for each $w \in \mathbb{D}$ the function

$$(1.1) \quad k_w^\alpha(z) = \frac{1 - \overline{\alpha(w)}\alpha(z)}{1 - \overline{w}z}, \quad z \in \mathbb{D},$$

2010 *Mathematics Subject Classification*: 47B32, 47B35, 30H10.

Key words and phrases: model spaces, truncated Toeplitz operators, asymmetric truncated Toeplitz operators, matrix representations.

December 12, 2016

is the corresponding kernel function, that is, $f(w) = \langle f, k_w^\alpha \rangle$ for every $f \in K_\alpha$ ($\langle \cdot, \cdot \rangle$ being the usual integral inner product). Since $k_w^\alpha \in H^\infty$, the set K_α^∞ of all bounded functions in K_α is dense in K_α .

It can be verified that for an inner function α the formula

$$(1.2) \quad C_\alpha f(z) = \alpha(z) \overline{zf(z)}, \quad |z| = 1,$$

defines an antilinear, isometric involution on $L^2(\partial\mathbb{D})$ and that the so-called conjugation C_α preserves K_α (see [18, Subection 2.3]). It is easy to see that the conjugate kernel $\tilde{k}_w^\alpha = C_\alpha k_w^\alpha$ is given by

$$\tilde{k}_w^\alpha(z) = \frac{\alpha(z) - \alpha(w)}{z - w}.$$

The unilateral shift is an example of a Toeplitz operator. The classical Toeplitz operator T_φ on H^2 with symbol $\varphi \in L^2(\partial\mathbb{D})$ is defined by

$$T_\varphi f = P(\varphi f).$$

The operator T_φ is densely defined on bounded functions and can be extended to a bounded operator on H^2 if and only if $\varphi \in L^\infty(\partial\mathbb{D})$. We have $S = T_z$.

In 2007 in his paper [18] D. Sarason initiated the study of the class compressions of classical Toeplitz operators to the model spaces, that is, the truncated Toeplitz operators.

Let α be an inner function and let P_α denote the orthogonal projection from $L^2(\partial\mathbb{D})$ onto K_α . A truncated Toeplitz operator on K_α with a symbol $\varphi \in L^2(\partial\mathbb{D})$ is given by

$$A_\varphi^\alpha f = P_\alpha(\varphi f).$$

The operator A_φ^α is defined on a dense subset K_α^∞ of K_α . Clearly, if $\varphi \in L^\infty(\partial\mathbb{D})$, then A_φ^α extends to a bounded operator on K_α . However, the boundedness of the symbol is not a necessary condition for the boundedness of the operator. Let $\mathcal{T}(\alpha)$ denote the set of all bounded truncated Toeplitz operators on K_α .

Among many interesting results concerning truncated Toeplitz operators (see [7, 12, 13, 14, 15]) there are the characterizations of these operators, given in [18] and [9].

It is known that the classical Toeplitz operators are those bounded linear operators $T: H^2 \rightarrow H^2$ that satisfy $T - S^*TS = 0$. In [18] Sarason gave a similar description of truncated Toeplitz operators on K_α using the compressed shift $S_\alpha = A_z^\alpha$. He proved that a bounded linear operator $A: K_\alpha \rightarrow K_\alpha$ belongs to $\mathcal{T}(\alpha)$ if and only if there exist $\chi, \psi \in K_\alpha$ such that

$$A - S_\alpha^* A S_\alpha = \psi \otimes \tilde{k}_0^\alpha + \tilde{k}_0^\alpha \otimes \chi$$

(here $f \otimes g$ is the standard rank-one operator, $f \otimes g(h) = \langle h, g \rangle f$). In fact, [18] contains a more general version of this result, where S_α is replaced by the so-called modified compressed shift. For $c \in \mathbb{C}$ the modified compressed shift $S_{\alpha,c}$ is given by

$$S_{\alpha,c} = S_\alpha + c(k_0^\alpha \otimes \tilde{k}_0^\alpha).$$

An operator A on K_α belongs to $\mathcal{T}(\alpha)$ if and only if for some $c \in \mathbb{C}$ there exist $\chi_c, \psi_c \in K_\alpha$ such that

$$A - S_{\alpha,c}^* A S_{\alpha,c} = \psi_c \otimes \tilde{k}_0^\alpha + \tilde{k}_0^\alpha \otimes \chi_c.$$

The equation $T - S^*TS = 0$ can be expressed in terms of the matrix representation of T with respect to the monomial basis of H^2 . It then states that T is a Toeplitz operator if and only if the matrix representing T is a Toeplitz matrix, i.e., an infinite matrix with constant diagonals. It follows that if $\alpha(z) = z^n$, then the operators from $\mathcal{T}(\alpha)$ are represented by finite Toeplitz matrices.

J. A. Cima, W. T. Ross and W.R. Wogen [9] considered the matrix representations of truncated Toeplitz operators in the case when K_α is a finite-dimensional model space, that is, when α is a finite Blaschke product. In particular, they proved that if α is a finite Blaschke product with distinct zeros $a_1, \dots, a_m$, then the matrix representing a truncated Toeplitz operator with respect to the reproducing kernel basis $\{k_{a_1}^\alpha, \dots, k_{a_m}^\alpha\}$ is completely determined by its entries along the main diagonal and the first row. They also gave similar characterizations in terms of matrix representations with respect to the so-called Clark bases and modified Clark bases (for detailed definitions see the following section).

Recently, in [4] and [6] the so-called asymmetric truncated Toeplitz operators were introduced. Let α, β be two inner functions. An asymmetric truncated Toeplitz operator $A_\varphi^{\alpha, \beta}$ on K_α with symbol $\varphi \in L^2(\partial\mathbb{D})$ is defined by

$$A_\varphi^{\alpha, \beta} f = P_\beta(\varphi f), \quad f \in K_\alpha.$$

Let

$$\mathcal{T}(\alpha, \beta) = \{A_\varphi^{\alpha, \beta} : \varphi \in L^2(\partial\mathbb{D}) \text{ and } A_\varphi^{\alpha, \beta} \text{ is bounded}\}.$$

Obviously, $A_\varphi^{\alpha, \alpha} = A_\varphi^\alpha$ and $\mathcal{T}(\alpha, \alpha) = \mathcal{T}(\alpha)$.

A characterization of operators from $\mathcal{T}(\alpha, \beta)$ in terms of compressed shifts was obtained in [4] for α and β such that β divides α (α/β is also an inner function), whereas the matrix representations of asymmetric truncated Toeplitz operators on finite-dimensional spaces were discussed in [17].

In this paper we continue the study of asymmetric truncated Toeplitz operators acting between finite-dimensional model spaces. In Section 2 we cite some results from [17] and then use these results in Section 3 to prove that the operators from $\mathcal{T}(\alpha, \beta)$ (in the finite-dimensional setting) can be characterized in terms of the modified compressed shifts $S_{\alpha, a}, S_{\beta, b}$. We also mention the characterization of asymmetric truncated Toeplitz operators in terms of the shift invariance.

In Section 4 we investigate the rank-one asymmetric truncated Toeplitz operators. We describe all the rank-one operators from $\mathcal{T}(\alpha, \beta)$ for finite Blaschke products α and β , and point out the cases when the given description differs from the one given by Sarason for rank-one truncated Toeplitz operators.

For the remainder of the paper we assume that the spaces K_α and K_β are finite-dimensional, that is, that α and β are two finite Blaschke products of degree m and n , respectively. Moreover, we assume that $m > 0$ and $n > 0$.

2. PRELIMINARIES

For any $\lambda_1 \in \partial\mathbb{D}$ define

$$(2.1) \quad U_{\lambda_1}^\alpha = S_\alpha + \frac{\lambda_1 + \alpha(0)}{1 - |\alpha(0)|^2} (k_0^\alpha \otimes \tilde{k}_0^\alpha).$$

Then $U_{\lambda_1}^\alpha : K_\alpha \rightarrow K_\alpha$ is unitary and, in fact, all one-dimensional unitary perturbations of the compressed shift S_α are of this form. This result is due to D.N. Clark [10] who also described the spectrum of $U_{\lambda_1}^\alpha$. In particular, he showed that the

set of eigenvalues of $U_{\lambda_1}^\alpha$ consists of those $\eta \in \partial\mathbb{D}$ at which α has a finite angular derivative and

$$(2.2) \quad \alpha(\eta) = \alpha_{\lambda_1} = \frac{\lambda_1 + \alpha(0)}{1 + \overline{\alpha(0)}\lambda_1}.$$

The corresponding eigenvectors are given by

$$k_\eta^\alpha(z) = \frac{1 - \overline{\alpha(\eta)}\alpha(z)}{1 - \overline{\eta}z}.$$

Actually, k_η^α belongs to K_α for every $\eta \in \partial\mathbb{D}$ such that α has an angular derivative in the sense of Carathéodory at η (α has an angular derivative at η and $|\alpha(\eta)| = 1$). In that case every $f \in K_\alpha$ has a non-tangential limit $f(\eta)$ at η and $f(\eta) = \langle f, k_\eta^\alpha \rangle$ (see [12, Thm. 7.4.1]). Note also that $\|k_\eta^\alpha\| = \sqrt{|\alpha'(\eta)|}$.

Here α is a finite Blaschke product, so k_η^α belongs to K_α for all $\eta \in \partial\mathbb{D}$. Since the degree of α is m , the equation (2.2) has precisely m distinct solutions $\eta_1, \dots, \eta_m$ on the unit circle $\partial\mathbb{D}$ (see [13, p. 6]). Moreover, since the dimension of K_α is m , the orthogonal set of corresponding eigenvectors $\{k_{\eta_1}^\alpha, \dots, k_{\eta_m}^\alpha\}$ forms a basis for K_α . The set of normalized kernel functions $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$,

$$v_{\eta_j}^\alpha = \|k_{\eta_j}^\alpha\|^{-1} k_{\eta_j}^\alpha, \quad j = 1, \dots, m,$$

is called the Clark basis corresponding to λ_1 (see [10] for more details).

Observe that for every $f \in K_\alpha$ we have

$$f = \sum_{j=1}^m \langle f, v_{\eta_j}^\alpha \rangle v_{\eta_j}^\alpha = \sum_{j=1}^m \frac{f(\eta_j)}{\sqrt{|\alpha'(\eta_j)|}} v_{\eta_j}^\alpha = \sum_{j=1}^m \frac{f(\eta_j)}{|\alpha'(\eta_j)|} k_{\eta_j}^\alpha.$$

If we put

$$e_{\eta_j}^\alpha = \omega_j^\alpha v_{\eta_j}^\alpha, \quad j = 1, \dots, m,$$

where

$$\omega_j^\alpha = e^{-\frac{i}{2}(\arg \eta_j - \arg \lambda_1)}, \quad j = 1, \dots, m,$$

then the basis $\{e_{\eta_1}^\alpha, \dots, e_{\eta_m}^\alpha\}$ has an additional property that

$$C_\alpha e_{\eta_j}^\alpha = e_{\eta_j}^\alpha, \quad j = 1, \dots, m,$$

where C_α is the conjugation given by (1.2). The basis $\{e_{\eta_1}^\alpha, \dots, e_{\eta_m}^\alpha\}$ is called the modified Clark basis.

Similarly, for $\lambda_2 \in \partial\mathbb{D}$ there are n orthogonal eigenvectors $k_{\zeta_1}^\beta, \dots, k_{\zeta_n}^\beta$ of the unitary operator

$$(2.3) \quad U_{\lambda_2}^\beta = S_\beta + \frac{\lambda_2 + \beta(0)}{1 - |\beta(0)|^2} (k_0^\beta \otimes \tilde{k}_0^\beta),$$

each corresponding to a different solution ζ_j , $j = 1, \dots, n$, of the equation

$$(2.4) \quad \beta(\zeta) = \beta_{\lambda_2} = \frac{\lambda_2 + \beta(0)}{1 + \overline{\beta(0)}\lambda_2}.$$

The Clark basis $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$ and modified Clark basis $\{e_{\zeta_1}^\beta, \dots, e_{\zeta_n}^\beta\}$ for K_β are defined analogously by

$$v_{\zeta_j}^\beta = \|k_{\zeta_j}^\beta\|^{-1} k_{\zeta_j}^\beta, \quad j = 1, \dots, n,$$

and

$$e_{\zeta_j}^\beta = \omega_j^\beta v_{\zeta_j}^\beta, \quad j = 1, \dots, n,$$

where

$$\omega_j^\beta = e^{-\frac{i}{2}(\arg \zeta_j - \arg \lambda_2)}, \quad j = 1, \dots, n.$$

Clearly, the equations (2.2) and (2.4) may have some solutions in common. In that case we assume that these solutions are precisely the numbers $\eta_j = \zeta_j$ for $j \leq l$ (with $l = 0$ if there are no solutions in common).

In what follows we will use the characterization of operators from $\mathcal{T}(\alpha, \beta)$ in terms of their matrix representations with respect to the Clark bases.

Theorem 2.1 ([17], Thm. 3.4). *Let α and β be two finite Blaschke products of degree $m > 0$ and $n > 0$, respectively. Let $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ be the Clark basis for K_α corresponding to $\lambda_1 \in \partial\mathbb{D}$, let $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$ be the Clark basis for K_β corresponding to $\lambda_2 \in \partial\mathbb{D}$ and assume that the sets $\{\eta_1, \dots, \eta_m\}$, $\{\zeta_1, \dots, \zeta_n\}$ have precisely l elements in common: $\eta_j = \zeta_j$ for $j \leq l$ ($l = 0$ if there are no elements in common). Finally, let A be any linear transformation from K_α into K_β . If $M_A = (r_{s,p})$ is the matrix representation of A with respect to the bases $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ and $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$, and*

(a) $l = 0$, then $A \in \mathcal{T}(\alpha, \beta)$ if and only if

$$(2.5) \quad r_{s,p} = \left(\frac{\sqrt{|\alpha'(\eta_1)|}}{\sqrt{|\alpha'(\eta_p)|}} \frac{\eta_p}{\eta_1} \frac{\eta_1 - \zeta_s}{\eta_p - \zeta_s} r_{s,1} + \frac{\sqrt{|\alpha'(\eta_1)|} \sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p}{\eta_1} \frac{\zeta_1 - \eta_1}{\eta_p - \zeta_s} r_{1,1} \right. \\ \left. + \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p - \zeta_1}{\eta_p - \zeta_s} r_{1,p} \right)$$

for all $1 \leq p \leq m$ and $1 \leq s \leq n$;

(b) $l > 0$, then $A \in \mathcal{T}(\alpha, \beta)$ if and only if

$$(2.6) \quad r_{s,p} = \left(\frac{\sqrt{|\alpha'(\eta_s)|} \sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p}{\eta_s} \frac{\eta_1 - \zeta_s}{\eta_p - \zeta_s} r_{1,s} + \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p - \zeta_1}{\eta_p - \zeta_s} r_{1,p} \right)$$

for all p, s such that $1 \leq p \leq m$, $1 \leq s \leq l$, $s \neq p$, and

$$(2.7) \quad r_{s,p} = \left(\frac{\sqrt{|\alpha'(\eta_1)|}}{\sqrt{|\alpha'(\eta_p)|}} \frac{\eta_p}{\eta_1} \frac{\eta_1 - \zeta_s}{\eta_p - \zeta_s} r_{s,1} + \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p - \zeta_1}{\eta_p - \zeta_s} r_{1,p} \right)$$

for all p, s such that $1 \leq p \leq m$, $l < s \leq n$.

3. CHARACTERIZATIONS USING RANK-TWO OPERATORS

In this section we prove that if α and β are finite Blaschke products, then a characterization in terms of modified compressed shifts, similar to that from [18, Thm. 10.1], can be given for operators from $\mathcal{T}(\alpha, \beta)$.

Theorem 3.1. *Let α, β be two finite Blaschke products and let a, b be two complex numbers. A linear operator A from K_α into K_β belongs to $\mathcal{T}(\alpha, \beta)$ if and only if there are functions $\chi_{a,b} \in K_\alpha$ and $\psi_{a,b} \in K_\beta$ such that*

$$(3.1) \quad A - S_{\beta,b} A S_{\alpha,a}^* = \psi_{a,b} \otimes k_0^\alpha + k_0^\beta \otimes \chi_{a,b}.$$

Proof. Let A be a linear operator from K_α into K_β and let a, b be two complex numbers.

Note first that if a_1, b_1 are any other two complex numbers, then the operator A satisfies (3.1) with some $\chi_{a,b} \in K_\alpha$, $\psi_{a,b} \in K_\beta$ if and only if there exist $\chi_{a_1,b_1} \in K_\alpha$, $\psi_{a_1,b_1} \in K_\beta$ such that

$$(3.2) \quad A - S_{\beta,b_1}AS_{\alpha,a_1}^* = \psi_{a_1,b_1} \otimes k_0^\alpha + k_0^\beta \otimes \chi_{a_1,b_1}.$$

Indeed, using

$$\begin{aligned} S_{\beta,b}AS_{\alpha,a}^* &= [S_\beta + b(k_0^\beta \otimes \tilde{k}_0^\beta)]A[S_\alpha^* + \bar{a}(\tilde{k}_0^\alpha \otimes k_0^\alpha)] \\ &= S_\beta AS_\alpha^* + b(k_0^\beta \otimes S_\alpha A^* \tilde{k}_0^\beta) + \bar{a}(S_\beta A \tilde{k}_0^\alpha \otimes k_0^\alpha) + b\bar{a}\langle A \tilde{k}_0^\alpha, \tilde{k}_0^\beta \rangle (k_0^\beta \otimes k_0^\alpha), \end{aligned}$$

and an analogous formula for $S_{\beta,b_1}AS_{\alpha,a_1}^*$, it is easy to verify that there exist functions $\chi \in K_\alpha$ and $\psi \in K_\beta$ such that

$$A - S_{\beta,b_1}AS_{\alpha,a_1}^* = A - S_{\beta,b}AS_{\alpha,a}^* + \psi \otimes k_0^\alpha + k_0^\beta \otimes \chi.$$

This implies the equivalence of (3.1) and (3.2).

In other words, instead of the condition (3.1) we can consider the condition (3.2) with a suitable a_1, b_1 .

Fix $\lambda_1, \lambda_2 \in \partial\mathbb{D}$ and let

$$a_1 = \frac{\alpha(0) + \lambda_1}{1 - |\alpha(0)|^2} \quad \text{and} \quad b_1 = \frac{\beta(0) + \lambda_2}{1 - |\beta(0)|^2}.$$

Since

$$S_{\alpha,a_1} = U_{\lambda_1}^\alpha \quad \text{and} \quad S_{\beta,b_1} = U_{\lambda_2}^\beta,$$

where $U_{\lambda_1}^\alpha$ and $U_{\lambda_2}^\beta$ are the Clark operators defined by (2.1) and (2.3), the condition (3.2) becomes

$$(3.3) \quad A - U_{\lambda_2}^\beta A(U_{\lambda_1}^\alpha)^* = \psi \otimes k_0^\alpha + k_0^\beta \otimes \chi,$$

with $\chi = \chi_{a_1,b_1} \in K_\alpha$ and $\psi = \psi_{a_1,b_1} \in K_\beta$.

Let $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ and $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$ be the Clark bases of K_α and K_β corresponding to λ_1 and λ_2 , respectively. Moreover, let $M_A = (r_{s,p})$ be the matrix representation of A with respect to these bases.

We now show that $A \in \mathcal{T}(\alpha, \beta)$ if and only if it satisfies (3.3). To this end we, express the condition (3.3) in terms of the matrix representation M_A and use Theorem 2.1.

The operators in (3.3) are equal if and only if their matrix representations with respect to $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ and $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$ are equal. The latter holds if and only if

$$(3.4) \quad \begin{aligned} \langle Av_{\eta_p}^\alpha, v_{\zeta_s}^\beta \rangle - \langle A(U_{\lambda_1}^\alpha)^* v_{\eta_p}^\alpha, (U_{\lambda_2}^\beta)^* v_{\zeta_s}^\beta \rangle \\ = \langle v_{\eta_p}^\alpha, k_0^\alpha \rangle \langle \psi, v_{\zeta_s}^\beta \rangle + \langle v_{\eta_p}^\alpha, \chi \rangle \langle k_0^\beta, v_{\zeta_s}^\beta \rangle \end{aligned} \quad \text{for all} \quad \begin{matrix} 1 \leq p \leq m \\ 1 \leq s \leq n \end{matrix}.$$

Recall that

$$r_{s,p} = \langle Av_{\eta_p}^\alpha, v_{\zeta_s}^\beta \rangle.$$

This, together with

$$(U_{\lambda_1}^\alpha)^* v_{\eta_p}^\alpha = \bar{\eta}_p v_{\eta_p}^\alpha, \quad (U_{\lambda_2}^\beta)^* v_{\zeta_s}^\beta = \bar{\zeta}_s v_{\zeta_s}^\beta,$$

and the reproducing property of the eigenvectors of $U_{\lambda_1}^\alpha$ and $U_{\lambda_2}^\beta$, implies that the system of equations (3.4) is equivalent to the following

$$(3.5) \quad \begin{aligned} (1 - \bar{\eta}_p \zeta_s) r_{s,p} \sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|} \\ = \psi(\zeta_s) \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_s) \overline{\chi(\eta_p)} \end{aligned} \quad \begin{aligned} 1 \leq p \leq m \\ 1 \leq s \leq n \end{aligned}.$$

Assume that A satisfies (3.3) with some $\chi \in K_\alpha$ and $\psi \in K_\beta$. By the above, the matrix representation $M_A = (r_{s,p})$ satisfies (3.5). If $l = 0$, then

$$\begin{aligned} \left(\frac{\sqrt{|\alpha'(\eta_1)|}}{\sqrt{|\alpha'(\eta_p)|}} \frac{\eta_p}{\eta_1} \frac{\eta_1 - \zeta_s}{\eta_p - \zeta_s} r_{s,1} + \frac{\sqrt{|\alpha'(\eta_1)|} \sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p}{\eta_1} \frac{\zeta_1 - \eta_1}{\eta_p - \zeta_s} r_{1,1} \right. \\ \left. + \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p - \zeta_1}{\eta_p - \zeta_s} r_{1,p} \right) = \frac{\psi(\zeta_s) \overline{k_0^\alpha(\eta_1)} + k_0^\beta(\zeta_1) \overline{\chi(\eta_p)}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|} (1 - \bar{\eta}_p \zeta_s)} \\ = \frac{\psi(\zeta_s) \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_s) \overline{\chi(\eta_p)}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|} (1 - \bar{\eta}_p \zeta_s)} = r_{s,p}, \end{aligned}$$

for all $1 \leq p \leq m$ and $1 \leq s \leq n$. This follows from (3.5) and the fact that

$$k_0^\alpha(\eta_p) = 1 - \overline{\alpha(0)} \alpha_{\lambda_1} = k_0^\alpha(\eta_1) \quad \text{and} \quad k_0^\beta(\zeta_s) = 1 - \overline{\beta(0)} \alpha_{\lambda_2} = k_0^\beta(\zeta_1).$$

By Theorem 2.1, A belongs to $\mathcal{T}(\alpha, \beta)$.

If $l > 0$, then (3.5) implies that

$$\psi(\zeta_s) \overline{k_0^\alpha(\eta_s)} + k_0^\beta(\zeta_s) \overline{\chi(\eta_s)} = 0$$

for all $1 \leq s \leq l$. It follows that

$$\begin{aligned} \left(\frac{\sqrt{|\alpha'(\eta_s)|}}{\sqrt{|\alpha'(\eta_p)|}} \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p}{\eta_s} \frac{\eta_1 - \zeta_s}{\eta_p - \zeta_s} r_{1,s} + \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p - \zeta_1}{\eta_p - \zeta_s} r_{1,p} \right) \\ = \frac{\psi(\zeta_1) \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_1) \overline{\chi(\eta_p)} - \psi(\zeta_1) \overline{k_0^\alpha(\eta_s)} - k_0^\beta(\zeta_1) \overline{\chi(\eta_s)}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|} (1 - \bar{\eta}_p \zeta_s)} = r_{s,p} \end{aligned}$$

for all $1 \leq p \leq m$, $1 \leq s \leq l$, $s \neq p$. Similarly,

$$\begin{aligned} \left(\frac{\sqrt{|\alpha'(\eta_1)|}}{\sqrt{|\alpha'(\eta_p)|}} \frac{\eta_p}{\eta_1} \frac{\eta_1 - \zeta_s}{\eta_p - \zeta_s} r_{s,1} + \frac{\sqrt{|\beta'(\zeta_1)|}}{\sqrt{|\beta'(\zeta_s)|}} \frac{\eta_p - \zeta_1}{\eta_p - \zeta_s} r_{1,p} \right) \\ = \frac{\psi(\zeta_s) \overline{k_0^\alpha(\eta_1)} + k_0^\beta(\zeta_s) \overline{\chi(\eta_1)} + \psi(\zeta_1) \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_1) \overline{\chi(\eta_p)}}{\sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|} (1 - \bar{\eta}_p \zeta_s)} = r_{s,p}. \end{aligned}$$

for all $1 \leq p \leq m$ and $l < s \leq n$. By Theorem 2.1 again, A belongs to $\mathcal{T}(\alpha, \beta)$.

Assume now that A belongs to $\mathcal{T}(\alpha, \beta)$. By the above, to show that there exist $\chi \in K_\alpha$ and $\psi \in K_\beta$ such that (3.3) holds it is enough to find finite sequences $\{\chi_1, \dots, \chi_m\}$ and $\{\psi_1, \dots, \psi_n\}$ such that

$$(3.6) \quad \psi_s \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_s) \overline{\chi_p} = (1 - \bar{\eta}_p \zeta_s) r_{s,p} \sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_s)|}, \quad \begin{aligned} 1 \leq p \leq m \\ 1 \leq s \leq n \end{aligned}.$$

Indeed, if $\{\chi_1, \dots, \chi_m\}$ and $\{\psi_1, \dots, \psi_n\}$ satisfy (3.6), then A satisfies (3.3) with χ and ψ given by

$$(3.7) \quad \chi = \sum_{p=1}^m \frac{\chi_p}{\sqrt{|\alpha'(\eta_p)|}} v_{\eta_p}^\alpha, \quad \psi = \sum_{s=1}^n \frac{\psi_s}{\sqrt{|\beta'(\zeta_s)|}} v_{\zeta_s}^\beta.$$

We now find such $\{\chi_1, \dots, \chi_m\}$ and $\{\psi_1, \dots, \psi_n\}$.

If $l = 0$, then, by Theorem 2.1(a), we only need to consider those equations from (3.6), which correspond to the elements $r_{1,p}$ and $r_{s,1}$ of the matrix representation of A . That is, $\{\chi_1, \dots, \chi_m\}$ and $\{\psi_1, \dots, \psi_n\}$ satisfy (3.6) if and only if they satisfy the following system of equations

$$(3.8) \quad \begin{cases} \psi_1 \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_1) \overline{\chi_p} = (1 - \overline{\eta_p} \zeta_1) r_{1,p} \sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_1)|} & \text{for all } 1 \leq p \leq m \\ \psi_s \overline{k_0^\alpha(\eta_1)} + k_0^\beta(\zeta_s) \overline{\chi_1} = (1 - \overline{\eta_1} \zeta_s) r_{s,1} \sqrt{|\alpha'(\eta_1)|} \sqrt{|\beta'(\zeta_s)|} & \text{for all } 1 < s \leq n \end{cases}.$$

This can easily be verified using (2.5).

Fix an arbitrary ψ_1 . Then the desired sequences are

$$\begin{cases} \chi_p = \frac{(1 - \overline{\eta_p} \zeta_1) \overline{r_{1,p}} \sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_1)|} - \overline{\psi_1} k_0^\alpha(\eta_p)}{k_0^\beta(\zeta_1)} & \text{for } 1 \leq p \leq m \\ \psi_s = \frac{(1 - \overline{\eta_1} \zeta_s) r_{s,1} \sqrt{|\alpha'(\eta_1)|} \sqrt{|\beta'(\zeta_s)|} - k_0^\beta(\zeta_s) \overline{\chi_1}}{k_0^\alpha(\eta_1)} & \text{for } 1 < s \leq n \end{cases},$$

and the functions χ and ψ in (3.3) are given by (3.7).

Similarly, if $l > 0$, then, by Theorem 2.1(b), $\{\chi_1, \dots, \chi_m\}$ and $\{\psi_1, \dots, \psi_n\}$ satisfy (3.6) if and only if they satisfy

$$(3.9) \quad \begin{cases} \psi_1 \overline{k_0^\alpha(\eta_p)} + k_0^\beta(\zeta_1) \overline{\chi_p} = (1 - \overline{\eta_p} \zeta_1) r_{1,p} \sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_1)|} & \text{for all } 1 \leq p \leq m \\ \psi_s \overline{k_0^\alpha(\eta_s)} + k_0^\beta(\zeta_s) \overline{\chi_s} = 0 & \text{for all } 1 \leq s \leq l \\ \psi_s \overline{k_0^\alpha(\eta_1)} + k_0^\beta(\zeta_s) \overline{\chi_1} = (1 - \overline{\eta_1} \zeta_s) r_{s,1} \sqrt{|\alpha'(\eta_1)|} \sqrt{|\beta'(\zeta_s)|} & \text{for all } l < s \leq n \end{cases}$$

(consider those equations from (3.6) with $r_{1,p}$ for $1 \leq p \leq m$, $r_{s,s}$ for $1 \leq s \leq l$ and $r_{s,1}$ for $l < s \leq n$). Fix an arbitrary ψ_1 . Then

$$\begin{cases} \chi_p = \frac{(1 - \overline{\eta_p} \zeta_1) \overline{r_{1,p}} \sqrt{|\alpha'(\eta_p)|} \sqrt{|\beta'(\zeta_1)|} - \overline{\psi_1} k_0^\alpha(\eta_p)}{k_0^\beta(\zeta_1)} & \text{for } 1 \leq p \leq m \\ \psi_s = -\frac{k_0^\beta(\zeta_s) \overline{\chi_s}}{k_0^\alpha(\eta_s)} & \text{for } 1 \leq s \leq l \\ \psi_s = \frac{(1 - \overline{\eta_1} \zeta_s) r_{s,1} \sqrt{|\alpha'(\eta_1)|} \sqrt{|\beta'(\zeta_s)|} - k_0^\beta(\zeta_s) \overline{\chi_1}}{k_0^\alpha(\eta_1)} & \text{for } l < s \leq n \end{cases}.$$

Again, A satisfies (3.3) with χ and ψ given by (3.7). This completes the proof. $\square$

Corollary 3.2. *Let α, β be two finite Blaschke products and let a, b be two complex numbers. A linear operator A from K_α into K_β belongs to $\mathcal{T}(\alpha, \beta)$ if and only if there are functions $\chi_{a,b} \in K_\alpha$ and $\psi_{a,b} \in K_\beta$ such that*

$$(3.10) \quad A - S_{\beta,b}^* A S_{\alpha,a} = \psi_{a,b} \otimes \tilde{k}_0^\alpha + \tilde{k}_0^\beta \otimes \chi_{a,b}.$$

Proof. We first prove that the linear operator A belongs to $\mathcal{T}(\alpha, \beta)$ if and only if $B = C_\beta A C_\alpha$ belongs to $\mathcal{T}(\alpha, \beta)$.

If A belongs to $\mathcal{T}(\alpha, \beta)$, then $A = A_\varphi^{\alpha,\beta}$ for some $\varphi \in L^2(\partial\mathbb{D})$, and for every $f \in K_\alpha^\infty$ and $g \in K_\beta^\infty$,

$$\begin{aligned} \langle Bf, g \rangle &= \langle C_\beta A C_\alpha f, g \rangle = \langle C_\beta g, A C_\alpha f \rangle \\ &= \langle \beta \overline{z} g, \varphi \alpha \overline{z} f \rangle = \langle \overline{\alpha \varphi} \beta f, g \rangle = \langle A_{\overline{\alpha \varphi} \beta}^{\alpha,\beta} f, g \rangle. \end{aligned}$$

Hence

$$B = C_\beta A C_\alpha = A_{\overline{\alpha \varphi} \beta}^{\alpha,\beta}.$$

On the other hand, if $B = C_\beta AC_\alpha$ belongs to $\mathcal{T}(\alpha, \beta)$, then, by the above, $C_\beta BC_\alpha$ also belongs to $\mathcal{T}(\alpha, \beta)$. However,

$$C_\beta BC_\alpha = C_\beta^2 AC_\alpha^2 = A,$$

and $A \in \mathcal{T}(\alpha, \beta)$.

We now prove that $A \in \mathcal{T}(\alpha, \beta)$ if and only if (3.10) holds.

Assume that $A \in \mathcal{T}(\alpha, \beta)$. Then $C_\beta AC_\alpha \in \mathcal{T}(\alpha, \beta)$ and, by Theorem 3.1, there are $\chi \in K_\alpha$ and $\psi \in K_\beta$ such that

$$(3.11) \quad C_\beta AC_\alpha - S_{\beta,b} C_\beta AC_\alpha S_{\alpha,a}^* = \psi \otimes k_0^\alpha + k_0^\beta \otimes \chi.$$

Since

$$C_\beta S_{\beta,b} C_\beta = S_{\beta,b}^* \quad \text{and} \quad C_\alpha S_{\alpha,a}^* C_\alpha = S_{\alpha,a},$$

we get

$$\begin{aligned} A - S_{\beta,b}^* A S_{\alpha,a} &= C_\beta^2 AC_\alpha^2 - C_\beta S_{\beta,b} C_\beta AC_\alpha S_{\alpha,a}^* C_\alpha \\ &= C_\beta (\psi \otimes k_0^\alpha + k_0^\beta \otimes \chi) C_\alpha = \tilde{\psi} \otimes \tilde{k}_0^\alpha + \tilde{k}_0^\beta \otimes \tilde{\chi}. \end{aligned}$$

Thus A satisfies (3.10) with

$$\chi_{a,b} = \tilde{\chi} \quad \text{and} \quad \psi_{a,b} = \tilde{\psi}.$$

A similar reasoning shows that if A satisfies (3.10), then $C_\beta AC_\alpha$ satisfies (3.1). By Theorem 3.1, $C_\beta AC_\alpha \in \mathcal{T}(\alpha, \beta)$ and so $A \in \mathcal{T}(\alpha, \beta)$. This completes the proof. $\square$

Corollary 3.3. *If α, β are two finite Blaschke products, then $\mathcal{T}(\alpha, \beta)$ is closed in the weak operator topology.*

Proof. The proof is analogous to that of [18, Thm. 4.2] and is thus left for the reader. $\square$

In [18, p. 512] D. Sarason considered the notion of shift invariance of operators on K_α . In [5] this notion was generalized to operators from K_α into K_β . A bounded linear operator from K_α into K_β is said to be shift invariant if

$$(3.12) \quad \langle ASf, Sg \rangle = \langle Af, g \rangle$$

for all $f \in K_\alpha$ and $g \in K_\beta$ such that $Sf \in K_\alpha$ and $Sg \in K_\beta$.

The authors in [5] proved that if α and β are two inner functions such that β divides α , then the set of all shift invariant operators from K_α into K_β is equal to $\mathcal{T}(\alpha, \beta)$. For finite Blaschke products we have the following.

Theorem 3.4. *Let α, β be two finite Blaschke products and let A be a linear operator from K_α into K_β . Then A belongs to $\mathcal{T}(\alpha, \beta)$ if and only if A is shift invariant.*

Proof. See [5]. $\square$

4. RANK-ONE ASYMMETRIC TRUNCATED TOEPLITZ OPERATORS

In Section 5 of [18] D. Sarason characterized all the truncated Toeplitz operators of rank-one. He proved (see [18, Thm. 5.1]) that every rank-one operator form $\mathcal{T}(\alpha)$ is a scalar multiple of $\tilde{k}_w^\alpha \otimes k_w^\alpha$ or $k_w^\alpha \otimes \tilde{k}_w^\alpha$ for some $w \in \overline{\mathbb{D}}$. Can a similar characterization be given for asymmetric truncated Toeplitz operators of rank-one?

It is known (see for instance [16, Prop. 3.1]) that the operators $\tilde{k}_w^\beta \otimes k_w^\alpha$ and $k_w^\beta \otimes \tilde{k}_w^\alpha$ belong to $\mathcal{T}(\alpha, \beta)$ for all $w \in \mathbb{D}$ and for such $w \in \partial\mathbb{D}$ that α and β have an angular derivative in the sense of Carathéodory at w . So a natural question is: is every rank-one operator from $\mathcal{T}(\alpha, \beta)$ a scalar multiple of one of these? A simple example shows that this is not always the case.

Example 4.1. Fix $a \in \mathbb{D}$, $a \neq 0$, and let

$$\alpha(z) = -z \frac{a-z}{1-\bar{a}z} \frac{a+z}{1+\bar{a}z}, \quad \beta(z) = z.$$

Then β divides α , K_β is a one-dimensional linear space spanned by $k_0^\beta = 1$ and K_α is a three-dimensional linear space spanned by $k_0^\alpha = 1$, $k_a^\alpha = k_a$ and $k_{-a}^\alpha = k_{-a}$. Moreover, since α and β are analytic in a domain containing $\overline{\mathbb{D}}$, the operators $\tilde{k}_w^\beta \otimes k_w^\alpha$ and $k_w^\beta \otimes \tilde{k}_w^\alpha$ belong to $\mathcal{T}(\alpha, \beta)$ for all $w \in \overline{\mathbb{D}}$.

It is easy to verify that $A = 1 \otimes (1 + k_a)$ is an element of $\mathcal{T}(\alpha, \beta)$. In fact,

$$A = 1 \otimes (1 + k_a) = A_{1+\tilde{k}_a}^{\alpha, \beta}.$$

We show that there is no $w \in \overline{\mathbb{D}}$ such that A is a scalar multiple of $\tilde{k}_w^\beta \otimes k_w^\alpha$ or $k_w^\beta \otimes \tilde{k}_w^\alpha$.

Assume first that

$$(4.1) \quad 1 \otimes (1 + k_a) = c(\tilde{k}_w^\beta \otimes k_w^\alpha)$$

for some $w \in \overline{\mathbb{D}}$ and $c \neq 0$. Since $k_w^\beta = \tilde{k}_w^\beta = 1$, the above implies that

$$1 + k_a = \bar{c}k_w^\alpha.$$

Equivalently,

$$(4.2) \quad \begin{cases} \langle 1 + k_a, 1 \rangle &= \langle \bar{c}k_w^\alpha, 1 \rangle \\ \langle 1 + k_a, k_a \rangle &= \langle \bar{c}k_w^\alpha, k_a \rangle \\ \langle 1 + k_a, k_{-a} \rangle &= \langle \bar{c}k_w^\alpha, k_{-a} \rangle \end{cases}.$$

The first equation in (4.2) implies that $c = 2$. By the second equation in (4.2),

$$1 + \frac{1}{1 - |a|^2} = \frac{2}{1 - \bar{w}a},$$

and

$$w = \frac{a}{2 - |a|^2}.$$

However, by the third equation,

$$1 + \frac{1}{1 + |a|^2} = \frac{2}{1 + \bar{w}a},$$

and

$$w = \frac{a}{2 + |a|^2}.$$

Clearly, $\frac{a}{2-|a|^2} = \frac{a}{2+|a|^2}$ if and only if $a = 0$, and thus there is no $w \in \overline{\mathbb{D}}$ for which (4.1) holds.

Assume then that

$$(4.3) \quad 1 \otimes (1 + k_a) = c(k_w^\beta \otimes \tilde{k}_w^\alpha)$$

for some $w \in \overline{\mathbb{D}}$ and $c \neq 0$. As before, we must have

$$1 + k_a = \bar{c}\tilde{k}_w^\alpha,$$

which is equivalent to the following system of equations

$$(4.4) \quad \begin{cases} \langle 1 + k_a, 1 \rangle &= \langle \bar{c}\tilde{k}_w^\alpha, 1 \rangle \\ \langle 1 + k_a, k_a \rangle &= \langle \bar{c}\tilde{k}_w^\alpha, k_a \rangle \\ \langle 1 + k_a, k_{-a} \rangle &= \langle \bar{c}\tilde{k}_w^\alpha, k_{-a} \rangle \end{cases}.$$

Since $\langle \tilde{k}_w^\alpha, 1 \rangle = \alpha(w)/w$, the first equation in (4.4) gives

$$(4.5) \quad 2 = -\bar{c} \frac{a-w}{1-\bar{a}w} \frac{a+w}{1+\bar{a}w}.$$

Similarly,

$$\langle \tilde{k}_w^\alpha, k_a \rangle = \frac{\alpha(w)}{w-a}, \quad \langle \tilde{k}_w^\alpha, k_{-a} \rangle = \frac{\alpha(w)}{w+a},$$

and the second and third equation in (4.4) yield

$$(4.6) \quad 1 + \frac{1}{1-|a|^2} = \bar{c} \frac{w}{1-\bar{a}w} \frac{a+w}{1+\bar{a}w},$$

and

$$(4.7) \quad 1 + \frac{1}{1+|a|^2} = -\bar{c} \frac{w}{1+\bar{a}w} \frac{a-w}{1-\bar{a}w},$$

respectively. By (4.5) and (4.6),

$$w = \frac{2-|a|^2}{\bar{a}}.$$

However, by (4.5) and (4.7),

$$w = \frac{2+|a|^2}{\bar{a}}.$$

As before, there is no $w \in \overline{\mathbb{D}}$ for which (4.3) holds.

Note that in the above example every linear operator from K_α into K_β is a rank-one operator and belongs to $\mathcal{T}(\alpha, \beta)$. This happens for all finite Blaschke products α, β of degree m, n , respectively, and such that $m = 1$ or $n = 1$. Indeed, if α and β are finite Blaschke products of degree m and n , respectively, then the dimension of $\mathcal{T}(\alpha, \beta)$ is $m + n - 1$ (see [17, Prop. 2.1]). It follows that if $m = 1$ or $n = 1$, then

$$\dim \mathcal{T}(\alpha, \beta) = m + n - 1 = mn.$$

Since mn is the dimension of $\mathcal{L}(\alpha, \beta)$, the space of all linear operators from K_α into K_β , we get $\mathcal{T}(\alpha, \beta) = \mathcal{L}(\alpha, \beta)$.

So if one of the spaces K_α, K_β is one-dimensional, then the set of rank-one asymmetric truncated Toeplitz operators is "quite big". Is it ever "big enough" to contain operators which are not scalar multiples of $\tilde{k}_w^\beta \otimes k_w^\alpha$ or $k_w^\beta \otimes \tilde{k}_w^\alpha$? To answer this we need the following.

Proposition 4.2. *Let α be a finite Blaschke product of degree $m > 0$. Every $f \in K_\alpha$ is a scalar multiple of a reproducing kernel or a conjugate kernel if and only if $m \leq 2$.*

Proof. Let $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ be the Clark basis of K_α , corresponding to a fixed $\lambda \in \partial\mathbb{D}$.

We first prove that if every $f \in K_\alpha$ is a scalar multiple of a reproducing kernel or a conjugate kernel, then $m \leq 2$. To this end, we show that if $m > 2$, then there exists $f \in K_\alpha$ that is neither a scalar multiple of a reproducing kernel nor a scalar multiple of a conjugate kernel.

Observe, that if

$$k_w^\alpha = c_1 v_{\eta_1}^\alpha + \dots + c_m v_{\eta_m}^\alpha$$

for some $w \in \overline{\mathbb{D}}$, then

$$\begin{aligned} c_j &= \langle k_w^\alpha, v_{\eta_j}^\alpha \rangle = \|k_{\eta_j}^\alpha\|^{-1} k_w^\alpha(\eta_j) \\ &= \frac{1}{\sqrt{|\alpha'(\eta_j)|}} \frac{1 - \overline{\alpha(w)}\alpha(\eta_j)}{1 - \overline{w}\eta_j} = \frac{1}{\sqrt{|\alpha'(\eta_j)|}} \frac{1 - \overline{\alpha(w)}\alpha_\lambda}{1 - \overline{w}\eta_j}. \end{aligned}$$

Hence, if $w \notin \{\eta_1, \dots, \eta_m\}$, then $\alpha(w) \neq \alpha_\lambda$ and $c_j \neq 0$ for all $1 \leq j \leq m$. On the other hand, if $w = \eta_{j_0}$ for some $1 \leq j_0 \leq m$, then

$$(4.8) \quad c_j = \begin{cases} 0 & \text{for } j \neq j_0 \\ \frac{1}{\sqrt{|\alpha'(\eta_{j_0})|}} & \text{for } j = j_0 \end{cases}.$$

Similarly, if

$$\tilde{k}_w^\alpha = c_1 v_{\eta_1}^\alpha + \dots + c_m v_{\eta_m}^\alpha$$

for some $w \in \overline{\mathbb{D}}$, then

$$c_j = \langle \tilde{k}_w^\alpha, v_{\eta_j}^\alpha \rangle = \frac{1}{\sqrt{|\alpha'(\eta_j)|}} \frac{\alpha_\lambda - \alpha(w)}{\eta_j - w},$$

and $c_j \neq 0$ for all $1 \leq j \leq m$ if $w \notin \{\eta_1, \dots, \eta_m\}$, or c_j 's satisfy (4.8) if $w = \eta_{j_0}$ for some $1 \leq j_0 \leq m$.

Consider a linear combination

$$f = c_1 v_{\eta_1}^\alpha + \dots + c_m v_{\eta_m}^\alpha.$$

By the above, if f is a scalar multiple of k_w^α or $\tilde{k}_w^\alpha$, then all the coefficients c_j are nonzero or there is precisely one nonzero coefficient c_{j_0} . If $m > 2$ and f is a linear combination as above with at least one coefficient equal to zero and at least two nonzero coefficients (for example $c_1 = c_2 = 1$ and $c_j = 0$ for $j > 2$), then f is neither a scalar multiple of a reproducing kernel nor a scalar multiple of a conjugate kernel.

To complete the proof we show that if $m \leq 2$, then every $f \in K_\alpha$ is a scalar multiple of a reproducing kernel or a conjugate kernel. This is clear for $m = 1$.

Let $m = 2$ and take $f = c_1 v_{\eta_1}^\alpha + c_2 v_{\eta_2}^\alpha$. If $c_1 c_2 = 0$, then f is a scalar multiple of a reproducing kernel. Assume that $c_1 c_2 \neq 0$. We find $w \in \overline{\mathbb{D}}$ such that f is a scalar multiple of k_w^α or $\tilde{k}_w^\alpha$.

If $f = c k_w^\alpha$, then

$$\begin{cases} c_1 = \frac{c}{\sqrt{|\alpha'(\eta_1)|}} \frac{1 - \overline{\alpha(w)}\alpha_\lambda}{1 - \overline{w}\eta_1} \\ c_2 = \frac{c}{\sqrt{|\alpha'(\eta_2)|}} \frac{1 - \overline{\alpha(w)}\alpha_\lambda}{1 - \overline{w}\eta_2} \end{cases}.$$

From this

$$\frac{1 - \overline{w}\eta_1}{1 - \overline{w}\eta_2} = \frac{\sqrt{|\alpha'(\eta_2)|}}{\sqrt{|\alpha'(\eta_1)|}} \frac{c_2}{c_1},$$

and

$$(4.9) \quad w(\overline{\eta}_1 - \overline{w}_0\overline{\eta}_2) = 1 - \overline{w}_0,$$

where $w_0 = \frac{\sqrt{|\alpha'(\eta_2)|}}{\sqrt{|\alpha'(\eta_1)|}} \frac{c_2}{c_1}$.

On the other hand, if $f = c_1 v_{\eta_1}^\alpha + c_2 v_{\eta_2}^\alpha = \widetilde{c} k_w^\alpha$, then

$$\overline{c}_1 \alpha_\lambda \overline{\eta}_1 v_{\eta_1}^\alpha + \overline{c}_2 \alpha_\lambda \overline{\eta}_2 v_{\eta_2}^\alpha = \overline{c} k_w^\alpha.$$

As before,

$$\begin{cases} \overline{c}_1 \alpha_\lambda \overline{\eta}_1 = \frac{\overline{c}}{\sqrt{|\alpha'(\eta_1)|}} \frac{1 - \overline{\alpha(w)} \alpha_\lambda}{1 - \overline{w}\eta_1} \\ \overline{c}_2 \alpha_\lambda \overline{\eta}_2 = \frac{\overline{c}}{\sqrt{|\alpha'(\eta_2)|}} \frac{1 - \overline{\alpha(w)} \alpha_\lambda}{1 - \overline{w}\eta_2} \end{cases}$$

and

$$\frac{\overline{\eta}_1 - \overline{w}}{\overline{\eta}_2 - \overline{w}} = \frac{\sqrt{|\alpha'(\eta_2)|}}{\sqrt{|\alpha'(\eta_1)|}} \frac{\overline{c}_2}{\overline{c}_1} = \overline{w}_0.$$

From this

$$(4.10) \quad w(1 - w_0) = \eta_1 - w_0 \eta_2.$$

Therefore, we must show that there exists $w \in \overline{\mathbb{D}}$ such that it satisfies (4.9) (then f is a scalar multiple of k_w^α) or such that it satisfies (4.10) (then f is a scalar multiple of $\widetilde{k}_w^\alpha$). We consider three cases.

Case 1. $1 - w_0 = 0$.

In this case $w = 0$ satisfies (4.9), $c_1 \sqrt{|\alpha'(\eta_1)|} = c_2 \sqrt{|\alpha'(\eta_2)|}$ and $f = c k_0^\alpha$ with

$$c = \frac{c_1 \sqrt{|\alpha'(\eta_1)|}}{1 - \alpha(0) \alpha_\lambda} = \frac{c_2 \sqrt{|\alpha'(\eta_2)|}}{1 - \alpha(0) \alpha_\lambda}.$$

Case 2. $\eta_1 - w_0 \eta_2 = 0$.

In this case $w = 0$ satisfies (4.10), $c_1 \eta_1 \sqrt{|\alpha'(\eta_1)|} = c_2 \eta_2 \sqrt{|\alpha'(\eta_2)|}$ and $f = \widetilde{c} k_0^\alpha$ with

$$c = \frac{c_1 \eta_1 \sqrt{|\alpha'(\eta_1)|}}{\alpha_\lambda - \alpha(0)} = \frac{c_2 \eta_2 \sqrt{|\alpha'(\eta_2)|}}{\alpha_\lambda - \alpha(0)}.$$

Case 3. $1 - w_0 \neq 0$ and $\eta_1 - w_0 \eta_2 \neq 0$.

In this case (4.9) has a solution given by

$$w_1 = \frac{1 - \overline{w}_0}{\overline{\eta}_1 - \overline{w}_0 \overline{\eta}_2}$$

and (4.10) has a solution given by

$$w_2 = \frac{\eta_1 - w_0 \eta_2}{1 - w_0}.$$

Note that $w_2 = 1/\overline{w}_1$. Therefore, one of the numbers w_1, w_2 must be in $\overline{\mathbb{D}}$.

If $w_1 \in \overline{\mathbb{D}}$, then $f = c k_{w_1}^\alpha$ with

$$c = c_1 \sqrt{|\alpha'(\eta_1)|} \frac{1 - \overline{w}_1 \eta_1}{1 - \alpha(w_1) \alpha_\lambda} = c_2 \sqrt{|\alpha'(\eta_2)|} \frac{1 - \overline{w}_1 \eta_2}{1 - \alpha(w_1) \alpha_\lambda}.$$

If $w_2 \in \overline{\mathbb{D}}$, then $f = c\tilde{k}_{w_2}^\alpha$ with

$$c = c_1 \sqrt{|\alpha'(\eta_1)|} \frac{\eta_1 - w_2}{\alpha_\lambda - \alpha(w_2)} = c_2 \sqrt{|\alpha'(\eta_2)|} \frac{\eta_2 - w_2}{\alpha_\lambda - \alpha(w_2)}.$$

□

Corollary 4.3. *Let α and β be two finite Blaschke products of degree $m > 0$ and $n > 0$, respectively.*

- (a) *If $mn \leq 2$, then every operator from $\mathcal{T}(\alpha, \beta)$ is a scalar multiple of $\tilde{k}_w^\beta \otimes k_w^\alpha$ or $k_w^\beta \otimes \tilde{k}_w^\alpha$ for some $w \in \overline{\mathbb{D}}$.*
- (b) *If either $m = 1$ and $n > 2$, or $m > 2$ and $n = 1$, then there exists a rank-one operator from $\mathcal{T}(\alpha, \beta)$ that is neither a scalar multiple of $\tilde{k}_w^\beta \otimes k_w^\alpha$ nor a scalar multiple of $k_w^\beta \otimes \tilde{k}_w^\alpha$.*

Proof. (a) Let $f \in K_\alpha$ and $g \in K_\beta$ be such that $g \otimes f \in \mathcal{T}(\alpha, \beta)$.

If $m = n = 1$, then $f = c_1 k_0^\alpha$, $g = c_2 \tilde{k}_0^\beta$ and $g \otimes f$ is a scalar multiple of $\tilde{k}_0^\beta \otimes k_0^\alpha$.

If $m = 2$, $n = 1$, then, by Proposition 4.2, there exists $w \in \overline{\mathbb{D}}$ such that f is a scalar multiple of k_w^α or $\tilde{k}_w^\alpha$. Since here K_β is one-dimensional, g is always a scalar multiple of $\tilde{k}_w^\beta$ and of k_w^β . Hence $g \otimes f$ is a scalar multiple of $\tilde{k}_w^\beta \otimes k_w^\alpha$ or $k_w^\beta \otimes \tilde{k}_w^\alpha$.

A similar reasoning can be used for $m = 1$, $n = 2$.

(b) Let $m = 1$ and $n > 2$. By Proposition 4.2, there exists $g \in K_\beta$ that g is neither a scalar multiple of a reproducing kernel nor a scalar multiple of a conjugate kernel. The desired operator is $A = g \otimes f$ with any $f \in K_\alpha$, $f \neq 0$.

The proof for $m > 2$, $n = 1$ is similar. □

What happens if both m and n are greater than one? It turns out that in that case every rank-one operator from $\mathcal{T}(\alpha, \beta)$ is a scalar multiple of $\tilde{k}_w^\beta \otimes k_w^\alpha$ or $k_w^\beta \otimes \tilde{k}_w^\alpha$ for some $w \in \overline{\mathbb{D}}$. To give the proof we need the following.

Lemma 4.4. *Let α and β be two finite Blaschke products of degree $m > 1$ and $n > 1$, respectively. Let $f \in K_\alpha$, $g \in K_\beta$ be two nonzero functions such that $g \otimes f$ belongs to $\mathcal{T}(\alpha, \beta)$ and let $w \in \overline{\mathbb{D}}$. Then*

- (a) *g is a scalar multiple of k_w^β if and only if f is a scalar multiple of $\tilde{k}_w^\alpha$,*
- (b) *g is a scalar multiple of $\tilde{k}_w^\beta$ if and only if f is a scalar multiple of k_w^α .*

Proof. Fix $\lambda_1, \lambda_2 \in \partial\mathbb{D}$. Let $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ be the Clark basis for K_α corresponding to λ_1 and let $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$ be the Clark basis for K_β corresponding to λ_2 . Assume moreover that the sets $\{\eta_1, \dots, \eta_m\}$, $\{\zeta_1, \dots, \zeta_n\}$ have precisely l elements in common, these elements being $\eta_j = \zeta_j$, $j \leq l$.

Assume that $f \in K_\alpha$, $g \in K_\beta$ are such that $g \otimes f$ belongs to $\mathcal{T}(\alpha, \beta)$ and let $(r_{s,p})$ be the matrix representation of $g \otimes f$ with respect to $\{v_{\eta_1}^\alpha, \dots, v_{\eta_m}^\alpha\}$ and $\{v_{\zeta_1}^\beta, \dots, v_{\zeta_n}^\beta\}$.

We only show that if $g = k_w^\beta$, then f is a scalar multiple of $\tilde{k}_w^\alpha$, and that if $g = \tilde{k}_w^\beta$, then f is a scalar multiple of k_w^α . The proof in the general case is just a matter of multiplying by a constant. Moreover, the other implications then follow, because $g \otimes f$ belongs to $\mathcal{T}(\alpha, \beta)$ if and only if $f \otimes g$ belongs to $\mathcal{T}(\beta, \alpha)$.

We first give the proof for $l = 0$, that is, when $\{\eta_1, \dots, \eta_m\}$ and $\{\zeta_1, \dots, \zeta_n\}$ have no elements in common.

By Theorem 2.1, the matrix representation $(r_{s,p})$ satisfies (2.5). Since

$$r_{s,p} = \langle (g \otimes f)v_{\eta_p}^\alpha, v_{\zeta_s}^\beta \rangle = \langle g, v_{\zeta_s}^\beta \rangle \langle v_{\eta_p}^\alpha, f \rangle = \frac{g(\zeta_s)\overline{f(\eta_p)}}{\sqrt{|\beta'(\zeta_s)|}\sqrt{|\alpha'(\eta_p)|}},$$

the equation (2.5) can be written as

$$g(\zeta_s)\overline{f(\eta_p)} = \frac{1 - \bar{\eta}_1\zeta_s}{1 - \bar{\eta}_p\zeta_s}g(\zeta_s)\overline{f(\eta_1)} - \frac{1 - \bar{\eta}_1\zeta_1}{1 - \bar{\eta}_p\zeta_s}g(\zeta_1)\overline{f(\eta_1)} + \frac{1 - \bar{\eta}_p\zeta_1}{1 - \bar{\eta}_p\zeta_s}g(\zeta_1)\overline{f(\eta_p)},$$

or

$$(4.11) \quad [a_{s,p}g(\zeta_s) - a_{1,p}g(\zeta_1)]\overline{f(\eta_p)} = [a_{s,1}g(\zeta_s) - a_{1,1}g(\zeta_1)]\overline{f(\eta_1)},$$

where

$$a_{s,p} = 1 - \bar{\eta}_p\zeta_s, \quad 1 \leq p \leq m, \quad 1 \leq s \leq n.$$

Let $g = k_w^\beta$ with $w \in \mathbb{D}$. We need to consider three cases.

Case 1. $w = \zeta_{s_0}$ for some $1 \leq s_0 \leq n$.

In this case $g(\zeta_{s_0}) \neq 0$ and $g(\zeta_s) = 0$ for all $s \neq s_0$. If $s_0 > 1$, then (4.11) (with $s = s_0$) implies that

$$(4.12) \quad a_{s_0,p}g(\zeta_{s_0})\overline{f(\eta_p)} = a_{s_0,1}g(\zeta_{s_0})\overline{f(\eta_1)}$$

for all $1 \leq p \leq m$. If $s_0 = 1$, then (4.12) also follows from (4.11) (with $s = n$ for example). Since $a_{s_0,p} \neq 0$, we get

$$\begin{aligned} f(\eta_p) &= \frac{\bar{a}_{s_0,1}}{\bar{a}_{s_0,p}}f(\eta_1) = \frac{1 - \bar{\eta}_1\zeta_{s_0}}{1 - \bar{\eta}_p\zeta_{s_0}}f(\eta_1) \\ &= \frac{1 - \bar{\zeta}_{s_0}\eta_1}{1 - \alpha(\zeta_{s_0})\alpha_{\lambda_1}}f(\eta_1)k_{\zeta_{s_0}}^\alpha(\eta_p) = ck_{\zeta_{s_0}}^\alpha(\eta_p) \end{aligned}$$

for all $1 \leq p \leq m$. From this,

$$\langle f, v_{\eta_p}^\alpha \rangle = \langle ck_{\zeta_{s_0}}^\alpha, v_{\eta_p}^\alpha \rangle$$

for all $1 \leq p \leq m$ and f is a scalar multiple of $k_{\zeta_{s_0}}^\alpha$. Since

$$(4.13) \quad k_w^\alpha = \overline{\alpha(w)}w\tilde{k}_w^\alpha$$

for all $w \in \partial\mathbb{D}$, f is a scalar multiple of $\tilde{k}_{\zeta_{s_0}}^\alpha$.

Case 2. $w = \eta_{p_0}$ for some $1 \leq p_0 \leq m$.

Note that if $g = k_w^\beta$ with $w \in \mathbb{D} \setminus \{\zeta_1, \dots, \zeta_n\}$, then

$$(4.14) \quad \begin{aligned} a_{s,p}g(\zeta_s) - a_{1,p}g(\zeta_1) &= (1 - \bar{\eta}_p\zeta_s)\frac{1 - \overline{\beta(w)}\beta_{\lambda_2}}{1 - \bar{w}\zeta_s} - (1 - \bar{\eta}_p\zeta_1)\frac{1 - \overline{\beta(w)}\beta_{\lambda_2}}{1 - \bar{w}\zeta_1} \\ &= \frac{(1 - \overline{\beta(w)}\beta_{\lambda_2})(\bar{w} - \bar{\eta}_p)(\zeta_s - \zeta_1)}{(1 - \bar{w}\zeta_s)(1 - \bar{w}\zeta_1)}. \end{aligned}$$

It follows that in this case

$$(4.15) \quad a_{n,p_0}g(\zeta_n) - a_{1,p_0}g(\zeta_1) = 0,$$

and

$$(4.16) \quad a_{n,p}g(\zeta_n) - a_{1,p}g(\zeta_1) \neq 0$$

for all $p \neq p_0$.

If $p_0 = 1$, then (4.11) (with $s = n$) and (4.15) imply that

$$[a_{n,p}g(\zeta_n) - a_{1,p}g(\zeta_1)]\overline{f(\eta_p)} = [a_{n,1}g(\zeta_n) - a_{1,1}g(\zeta_1)]\overline{f(\eta_1)} = 0.$$

By (4.16), $f(\eta_p) = 0$ for all $p \neq p_0$. Clearly, f is a scalar multiple of $k_{\eta_{p_0}}^\alpha$ and so a scalar multiple of $\tilde{k}_{\eta_{p_0}}^\alpha$ by (4.13).

If $p_0 > 1$, then (4.11) (with $p = p_0$ and $s = n$) and (4.15) give

$$0 = [a_{n,p_0}g(\zeta_n) - a_{1,p_0}g(\zeta_1)]\overline{f(\eta_{p_0})} = [a_{n,1}g(\zeta_n) - a_{1,1}g(\zeta_1)]\overline{f(\eta_1)}.$$

By (4.16), $f(\eta_1) = 0$. From this and (4.11) again (with $s = n$),

$$[a_{n,p}g(\zeta_n) - a_{1,p}g(\zeta_1)]\overline{f(\eta_p)} = 0$$

for all p and therefore $f(\eta_p) = 0$ for all $p \neq p_0$. As before, f is a scalar multiple of $\tilde{k}_{\eta_{p_0}}^\alpha$.

Case 3. $w \notin \{\eta_1, \dots, \eta_m\} \cup \{\zeta_1, \dots, \zeta_n\}$.

In this case (4.11) (with $s = n$) gives

$$f(\eta_p) = \frac{\overline{a_{n,1}g(\zeta_n)} - \overline{a_{1,1}g(\zeta_1)}}{\overline{a_{n,p}g(\zeta_n)} - \overline{a_{1,p}g(\zeta_1)}} f(\eta_1)$$

for all $p > 1$. By (4.14),

$$\begin{aligned} f(\eta_p) &= \frac{w - \eta_1}{w - \eta_p} f(\eta_1) = \\ &= \frac{\eta_1 - w}{\alpha_{\lambda_1} - \alpha(w)} f(\eta_1) \tilde{k}_w^\alpha(\eta_p) = c \tilde{k}_w^\alpha(\eta_p). \end{aligned}$$

Hence f is a scalar multiple of $\tilde{k}_w^\alpha$.

This completes the proof of the first implication (for $l = 0$).

Now let $g = \tilde{k}_w^\beta$. If $w \in \partial\mathbb{D}$, then g is a scalar multiple of k_w^β by (4.13) and f is a scalar multiple of $\tilde{k}_w^\alpha$ by the first part of the proof. Assume that $w \in \mathbb{D}$. Then

$$(4.17) \quad a_{s,p}g(\zeta_s) - a_{1,p}g(\zeta_1) = \frac{(\beta_{\lambda_2} - \beta(w))(1 - \bar{\eta}_p w)(\zeta_1 - \zeta_s)}{(\zeta_s - w)(\zeta_1 - w)}.$$

By (4.11) (with $s = n$),

$$\begin{aligned} f(\eta_p) &= \frac{1 - \bar{w}\eta_1}{1 - \bar{w}\eta_p} f(\eta_1) \\ &= \frac{1 - \bar{w}\eta_1}{1 - \alpha(w)\alpha_{\lambda_1}} f(\eta_1) k_w^\alpha(\eta_p) = c k_w^\alpha(\eta_p) \end{aligned}$$

for all $1 \leq p \leq m$, and f is a scalar multiple of k_w^α .

This completes the proof of the second implication (for $l = 0$).

We now give the proof for $l > 0$. In this case the matrix representation $(r_{s,p})$ of $g \otimes f$ must satisfy (2.6) and (2.7). Instead of (4.11) we obtain

$$(4.18) \quad a_{s,p}g(\zeta_s)\overline{f(\eta_p)} = a_{1,p}g(\zeta_1)\overline{f(\eta_p)} - a_{1,s}g(\zeta_1)\overline{f(\eta_s)},$$

for all p, s such that $1 \leq p \leq m$, $1 \leq s \leq l$, $s \neq p$, and

$$(4.19) \quad a_{s,p}g(\zeta_s)\overline{f(\eta_p)} = a_{s,1}g(\zeta_s)\overline{f(\eta_1)} + a_{1,p}g(\zeta_1)\overline{f(\eta_p)},$$

for all p, s such that $1 \leq p \leq m$, $s > l$.

Let $g = k_w^\beta$ with $w \in \mathbb{D}$. As before, we have three cases to consider.

Case 1. $w = \zeta_{s_0}$ for some $1 \leq s_0 \leq n$.

In this case $g(\zeta_{s_0}) \neq 0$ and $g(\zeta_s) = 0$ for all $s \neq s_0$.

If $s_0 = 1 = l$, then (4.19) (with $s = n > 1$) gives

$$0 = a_{n,p}g(\zeta_n)\overline{f(\eta_p)} = a_{n,1}g(\zeta_n)\overline{f(\eta_1)} + a_{1,p}g(\zeta_1)\overline{f(\eta_p)} = a_{1,p}g(\zeta_1)\overline{f(\eta_p)}$$

for all $1 \leq p \leq m$ and so $f(\eta_p) = 0$ for all $p > 1$. Hence f is a scalar multiple of $k_{\eta_{s_0}}^\alpha = k_{\zeta_{s_0}}^\alpha$ (since here $\eta_{s_0} = \zeta_{s_0}$) and a scalar multiple of $\tilde{k}_{\zeta_{s_0}}^\alpha$ by (4.13).

If $s_0 = 1 < l$, then (4.18) (with $p = 1$) implies that

$$0 = a_{s,1}g(\zeta_s)\overline{f(\eta_1)} = -a_{1,s}g(\zeta_1)\overline{f(\eta_s)}$$

for every $s \neq s_0 = 1$, $s \leq l$. From this, $f(\eta_s) = 0$ for all $1 < s \leq l$. If $m = l$, then this implies that f is a scalar multiple of $k_{\zeta_{s_0}}^\alpha$ and as above, a scalar multiple of $\tilde{k}_{\zeta_{s_0}}^\alpha$. If $m > l$, then by (4.18) (with $s = l$, $p > l$),

$$0 = a_{l,p}g(\zeta_l)\overline{f(\eta_p)} = a_{1,p}g(\zeta_1)\overline{f(\eta_p)}$$

and $f(\eta_p) = 0$ for all $p > l$, which also implies that f is a scalar multiple of $\tilde{k}_{\zeta_{s_0}}^\alpha$.

If $1 < s_0 \leq l$, then (4.18) (with $s = s_0$) implies that

$$a_{s_0,p}g(\zeta_{s_0})\overline{f(\eta_p)} = 0$$

and $f(\eta_p) = 0$ for all $p \neq s_0$. Hence f is a scalar multiple of $\tilde{k}_{\zeta_{s_0}}^\alpha$.

If $1 \leq l < s_0$, then (4.19) (with $s = s_0$) gives

$$a_{s_0,p}g(\zeta_{s_0})\overline{f(\eta_p)} = a_{s_0,1}g(\zeta_{s_0})\overline{f(\eta_1)}$$

and

$$\begin{aligned} f(\eta_p) &= \frac{\overline{a_{s_0,1}}}{\overline{a_{s_0,p}}}f(\eta_1) = \frac{1 - \overline{\zeta_{s_0}}\eta_1}{1 - \overline{\zeta_{s_0}}\eta_p}f(\eta_1) \\ &= \frac{1 - \overline{\zeta_{s_0}}\eta_1}{1 - \alpha(\zeta_{s_0})\alpha_{\lambda_1}}f(\eta_1)k_{\zeta_{s_0}}^\alpha(\eta_p) = ck_{\zeta_{s_0}}^\alpha(\eta_p) \end{aligned}$$

for all $1 \leq p \leq m$, which again implies that f is a scalar multiple of $\tilde{k}_{\zeta_{s_0}}^\alpha$.

Case 2. $w = \eta_{p_0}$ for some $l < p_0 \leq m$.

In this case (4.14) implies that

$$a_{n,p_0}g(\zeta_n) - a_{1,p_0}g(\zeta_1) = 0,$$

and

$$a_{n,p}g(\zeta_n) - a_{1,p}g(\zeta_1) \neq 0$$

for all $p \neq p_0$.

If $l = n$, then (4.18) (with $p = p_0$, $s = n$) gives

$$0 = [a_{n,p_0}g(\zeta_n) - a_{1,p_0}g(\zeta_1)]\overline{f(\eta_{p_0})} = -a_{1,n}g(\zeta_1)\overline{f(\eta_n)}.$$

Since $g(\zeta_1) \neq 0$, we have $f(\eta_n) = 0$. By (4.18) again (with $s = n$),

$$(4.20) \quad [a_{n,p}g(\zeta_n) - a_{1,p}g(\zeta_1)]\overline{f(\eta_p)} = -a_{1,n}g(\zeta_1)\overline{f(\eta_n)} = 0$$

for $p \neq n$ and so $f(\eta_p) = 0$ for all $p \neq p_0$. Hence f is a scalar multiple of $\tilde{k}_{\eta_{p_0}}^\alpha$.

If $l < n$, then (4.19) (with $p = p_0$, $s = n$) implies that $f(\eta_1) = 0$ and that (4.20) holds for all $1 \leq p \leq m$. Hence $f(\eta_p) = 0$ for all $p \neq p_0$ and f is a scalar multiple of $\tilde{k}_{\eta_{p_0}}^\alpha$.

Case 3. $w \notin \{\eta_1, \dots, \eta_m\} \cup \{\zeta_1, \dots, \zeta_n\}$.

If $l = n$, then (4.18) (with $p = 1$, $s = n$) implies that

$$(4.21) \quad a_{n,1}g(\zeta_n)\overline{f(\eta_1)} = -a_{1,n}g(\zeta_1)\overline{f(\eta_n)}.$$

From this

$$\begin{aligned} f(\eta_n) &= -\frac{(1 - \eta_1\bar{\zeta}_n)(1 - w\bar{\zeta}_1)}{(1 - \eta_n\bar{\zeta}_1)(1 - w\bar{\zeta}_n)}f(\eta_1) \\ &= -\frac{(\zeta_n - \eta_1)(\zeta_1 - w)}{(\zeta_1 - \eta_n)(\zeta_n - w)}f(\eta_1) = \frac{\eta_1 - w}{\eta_n - w}f(\eta_1). \end{aligned}$$

Moreover, (4.18) (with $s = n$) together with (4.21) gives

$$(4.22) \quad [a_{n,p}g(\zeta_n) - a_{1,p}g(\zeta_1)]\overline{f(\eta_p)} = a_{n,1}g(\zeta_n)\overline{f(\eta_1)}$$

and, by (4.14),

$$\begin{aligned} f(\eta_p) &= \frac{\overline{a_{n,1}g(\zeta_n)}}{\overline{a_{n,p}g(\zeta_n)} - \overline{a_{1,p}g(\zeta_1)}}f(\eta_1) \\ &= \frac{(1 - \eta_1\bar{\zeta}_n)(1 - w\bar{\zeta}_1)}{(\zeta_1 - \bar{\zeta}_n)(\eta_p - w)}f(\eta_1) = \frac{\eta_1 - w}{\eta_p - w}f(\eta_1) \end{aligned}$$

for all $p \neq n$. Hence

$$f(\eta_p) = \frac{\eta_1 - w}{\eta_p - w}f(\eta_1) = \frac{\eta_1 - w}{\alpha_{\lambda_1} - \alpha(w)}f(\eta_1)\tilde{k}_w^\alpha(\eta_p) = c\tilde{k}_w^\alpha(\eta_p)$$

for all p , and f is a scalar multiple of $\tilde{k}_w^\alpha$.

If $l < n$, then it follows from (4.19) (with $s = n$) that (4.22) holds for all p . As above

$$f(\eta_p) = \frac{\eta_1 - w}{\eta_p - w}f(\eta_1) = c\tilde{k}_w^\alpha(\eta_p)$$

and f is a scalar multiple of $\tilde{k}_w^\alpha$.

This completes the proof of the first implication (for $l > 0$).

Now let $g = \tilde{k}_w^\beta$. As in the proof for $l = 0$, we can assume that $w \in \mathbb{D}$. Repeating the argument from Case 3 above one can show that

$$\begin{aligned} f(\eta_p) &= \frac{1 - \bar{w}\eta_1}{1 - \bar{w}\eta_p}f(\eta_1) \\ &= \frac{1 - \bar{w}\eta_1}{1 - \alpha(w)\alpha_{\lambda_1}}f(\eta_1)k_w^\alpha(\eta_p) = ck_w^\alpha(\eta_p) \end{aligned}$$

for all p . Hence f is a scalar multiple of k_w^α .

This completes the proof of the second implication (for $l > 0$) and the proof of the theorem. $\square$

Theorem 4.5. *Let α and β be two finite Blaschke products of degree $m > 0$ and $n > 0$, respectively. The only rank-one operators in $\mathcal{T}(\alpha, \beta)$ are the nonzero scalar multiples of the operators $\tilde{k}_w^\beta \otimes k_w^\alpha$ and $k_w^\beta \otimes \tilde{k}_w^\alpha$, $w \in \mathbb{D}$, if and only if either $mn \leq 2$, or $m > 1$ and $n > 1$.*

Proof. By Corollary 4.3, if the only rank-one operators in $\mathcal{T}(\alpha, \beta)$ are the nonzero scalar multiples of $\tilde{k}_w^\beta \otimes k_w^\alpha$ and $k_w^\beta \otimes \tilde{k}_w^\alpha$, $w \in \overline{\mathbb{D}}$, then either $mn \leq 2$, or $m > 1$ and $n > 1$.

To complete the proof we only need to show that if $m > 1$ and $n > 1$, then every rank-one operator from $\mathcal{T}(\alpha, \beta)$ is a scalar multiple of $\tilde{k}_w^\beta \otimes k_w^\alpha$ or $k_w^\beta \otimes \tilde{k}_w^\alpha$ for some $w \in \overline{\mathbb{D}}$. The proof is similar to that of [18, Thm. 5.1(c)]. Sarason's proof is based on the fact that every truncated Toeplitz operator is complex symmetric. Here one must use Lemma 4.4 instead. The details are left to the reader. $\square$

REFERENCES

- [1] P. R. Ahern, D. N. Clark, *Radial limits and invariant subspaces*, Amer. J. Math. 92 (1970), 332–342.
- [2] P. R. Ahern, D. N. Clark, *Radial n -th derivatives of Blaschke products*, Math. Scand. 28 (1971), 189–201.
- [3] A. Baranov, I. Chalendar, E. Fricain, J. E. Mashreghi, D. Timotin, *Bounded symbols and reproducing kernel thesis for truncated Toeplitz operators*, J. Funct. Anal. 259 (2010), no. 10, 2673–2701.
- [4] C. Cămară, J. Jurasik, K. Kliś-Garlicka, M. Ptak, *Characterizations of asymmetric truncated Toeplitz operators*, arXiv:1607.03342.
- [5] C. Cămară, K. Kliś-Garlicka, M. Ptak, *Shift invariance and reflexivity of asymmetric truncated Toeplitz operators*, preprint.
- [6] M. C. Cămară, J. R. Partington, *Asymmetric truncated Toeplitz operators and Toeplitz operators with matrix symbol*, arXiv:1504.06446.
- [7] J. A. Cima, S. R. Garcia, W. T. Ross, W. R. Wogen, *Truncated Toeplitz operators: spatial isomorphism, unitary equivalence, and similarity*, Indiana Univ. Math. J. 59 (2010), no. 2, 595–620.
- [8] J. A. Cima, A. L. Matheson, W. T. Ross, *The Cauchy transform*, Mathematical Surveys and Monographs, vol. 125, American Mathematical Society, Providence, RI, 2006.
- [9] J. A. Cima, W. T. Ross, W. R. Wogen, *Truncated Toeplitz operators on finite dimensional spaces*, Operators and Matrices 2 (2008), no. 3, 357–369.
- [10] D. N. Clark, *One dimensional perturbations of restricted shifts*, J. Anal. Math. 25 (1972), 169–191.
- [11] R. B. Crofoot, *Multipliers between invariant subspaces of the backward shift*, Pacific J. Math. 166 (1994), no. 2, 225–246.
- [12] S. R. Garcia, J. E. Mashreghi, W. Ross, *Introduction to Model Spaces and their Operators*, Cambridge University Press, 2016.
- [13] S. R. Garcia, W. T. Ross, *A nonlinear extremal problem on the Hardy space*, Comput. Methods Funct. Theory 9 (2009), no. 2, 485–524.
- [14] S. R. Garcia, W. T. Ross, *The norm of truncated Toeplitz operator*, CRM Proceedings and Lecture Notes 51 (2010), 59–64.
- [15] S. R. Garcia, W. T. Ross, *Recent progress on truncated Toeplitz operators*, in: J. Mashreghi, E. Fricain (Eds.), *Blaschke products and their applications*, Fields Inst. Commun., 65, Springer, New York, 2013, pp. 275–319.
- [16] J. Jurasik, B. Łanucha, *Asymmetric truncated Toeplitz operators equal to the zero operator*, in press, arXiv:1609.04039.
- [17] J. Jurasik, B. Łanucha, *Asymmetric truncated Toeplitz operators on finite-dimensional spaces*, in press, arXiv:1611.00813.
- [18] D. Sarason, *Algebraic properties of truncated Toeplitz operators*, Operators and Matrices 1 (2007), no. 4, 491–526.

DEPARTMENT OF MATHEMATICS, MARIA CURIE-SKŁODOWSKA UNIVERSITY, MARIA CURIE-SKŁODOWSKA SQUARE 1, 20-031 LUBLIN, POLAND

E-mail address: bartosz.lanucha@poczta.umcs.lublin.pl